

Comparison of antibacterial effect of methanolic and hydro-alcoholic ziziphus spina-christi extract with 2.5% sodium hypochlorite on enterococcus faecalis: an in vitro study

Akam Saeidi ¹, Mahmoud Reza Hamidi ², Abolfazl Davoodabadi ³, Elham Mahmoudi ², Soraya Khafri ⁴, Zahra Nikouee Rad[✉]

1. Assistant Professor, Dental Materials Research Center, Health Research Institute, Department of Endodontics, Babol University of Medical Sciences, Babol, IR Iran.
2. Assistant Professor, Department of Endodontics, Faculty of Dentistry, Babol University of Medical Sciences, Babol, IR Iran.
3. Assistant Professor, Department of Microbiology, Faculty of Medicine, Babol University of Medical Sciences, Babol, IR Iran.
4. Assistant Professor, Department of Biostatistics and Epidemiology, Faculty of Medicine, Babol University of Medical Sciences, Babol, IR Iran.
5. Dental student, Student Research Committee, Babol University of Medical Sciences, Babol, IR Iran.

✉Corresponding Author: Zahra Nikouee Rad, Faculty of Dentistry, Babol University of Medical Sciences, Babol, IR Iran.

Email: nikouee96@yahoo.com

Tel: +989119625188

Received: 25 Sept 2017

Accepted: 17 Mar 2018

Abstract

Introduction: Root treatment will not be successful, without a proper root canal irrigation with less disadvantages and antibacterial effect. The aim of this study was to compare antimicrobial effect of cedar extract and 2.5%NaOCl on *E. faecalis*.

Materials & Methods: In disk diffusion test, a standard suspension of *E. faecalis* (ATCC 29212) was cultured on plate and different concentrations (0.05, 0.15, 0.25, 0.35, 0.45 g/ml) of methanolic or hydro-alcoholic extracts, 2.5% NaOCl and physiologic serum (as negative control) were infused on paper disks. The inhibition zone measured after 48 h. In microdilution test, serial dilution of methanolic and hydro-alcoholic extracts, 2.5% NaOCl and physiologic serum in 1:2 proportion was performed in Brain Heart Infusion (BHI) culture medium. Then, standard suspension of *E. faecalis* was added to each well of micro plate. Data were analyzed using ANOVA.

Results: Hydro-alcoholic and methanolic extracts had antibacterial effect on *E. faecalis*. Inhibition zone of 2.5% NaOCl was significantly higher than that of other extracts ($p<0.001$). In microdilution test, *E. faecalis* bacterium was sensitive to both hydro-alcoholic and methanolic extracts but it was more sensitive to 2.5% NaOCl.

Conclusion: Totally, 2.5% NaOCl had the highest antibacterial effect on *E. faecalis* followed by hydro-alcoholic and methanolic extracts. NaOCl is an effective irrigant in root treatment until the studies like this can find a good alternative for it.

Keywords: Enterococcus faecalis, Methanol, Sodium hypochlorite, Ziziphus

Citation for article: Saeidi A, Hamidi MR, Davoodabadi A, Mahmoudi E, Khafri S, Nikouee Rad Z. An antibacterial effect comparison of methanolic and hydro-alcoholic extracts of ziziphus spina-christi with 2.5% sodium hypochlorite on enterococcus faecalis: an in vitro study. Caspian J Dent Res 2018; 7: 37-42.

مقایسه اثر ضد باکتری عصاره متانولی و هیدروالکلی گیاه سدر بر انتروکوکوس فکالیس با هیپوکلریت سدیم ۲/۵٪ درصد: (مطالعه آزمایشگاهی)

آکام سعیدی، محمودرضا حمیدی، ابوالفضل داوود آبادی، الهام محمودی، ثریا خفری، زهرا نیکویی راد*

چکیده

مقدمه: بدون یک شوینده مناسب کانال با عوارض کمتر و اثر ضد میکروبی، درمان ریشه موفق نخواهد بود. هدف این مطالعه مقایسه اثر ضد میکروبی عصاره سدر و هیپوکلریت سدیم ۲/۵٪ روی انتروکوکوس فکالیس است.

مواد و روش ها: در روش انتشار دیسک، یک سوسپانسیون استاندارد از باکتری انتروکوکوس فکالیس (ATCC 29212) روی محیط کشت داده شد و غلظت های مختلف عصاره های متانولی و هیدروالکلی سدر (۰/۰۵، ۰/۱۵، ۰/۲۵، ۰/۳۵، ۰/۴۵ g/ml)، هیپوکلریت سدیم ۲/۵٪ و سرم فیزیولوژی، روی دیسک های کاغذی ریخته شدند. بعد از ۴۸ ساعت هاله عدم رشد باکتری اندازه گیری شد. در روش میکروداپلوشن، رقیق سازی سریال عصاره های متانولی و هیدروالکلی، هیپوکلریت سدیم ۲/۵٪ و سرم فیزیولوژی در نسبت ۱:۲ در محیط BHI انجام شد و سپس سوسپانسیون استاندارد باکتری به خانه های میکروپلیت اضافه گردید. آنالیزهای آماری با استفاده از آزمون ANOVA انجام شد.

یافته ها: عصاره های متانولی و هیدروالکلی اثر ضد میکروبی روی باکتری انتروکوکوس فکالیس داشتند. قطر هاله عدم رشد هیپوکلریت سدیم ۲/۵٪ در مقایسه با عصاره ها به طور معناداری بزرگتر بود ($p < 0.001$). در تست میکروداپلوشن، انتروکوکوس فکالیس به هر دو عصاره متانولی و هیدروالکلی حساس بود اما نسبت به هیپوکلریت سدیم ۲/۵٪ بیشتر حساس بود.

نتیجه گیری: به این ترتیب هیپوکلریت سدیم ۲/۵٪، عصاره های هیدروالکلی و متانولی بیشترین اثر ضد میکروبی را نشان دادند. هیپوکلریت سدیم یک شستشودهنده موثر در درمان ریشه است تا زمانی که مطالعاتی اینچنینی بتوانند جایگزین مناسبی برای آن پیدا کنند.

واژگان کلیدی: انتروکوکوس فکالیس، متانول، هیپوکلریت سدیم، Ziziphus

Introduction

Bacteria are the most important factor in creating pulp and periapical disease. When trauma and caries cause pathologic changes in pulp of tooth, microorganisms get into pulp system and invade root canal. Because of bacteria competition for reaching to low amount of food and oxygen in root canal, specific number of bacteria species in endodontic infections will persist. *E. faecalis* is a facultative anaerobe Gram-positive bacterium which is one of the bacteria in constant periapical lesions. It can attack dentinal tubules and cope with hard condition of root canal, so it is a resistant microorganism.^[1]

Effective cleaning and complete disinfection of root canal system are needed to achieve long-term success of root canal therapy.^[2] Root canal cleaning contains two parts: instrumentation and irrigation processes. Instrumentation removes pulp remains, scrapes dentin from root walls and prepares them for aggression from canal. Irrigators remove debris of filing process from root canal and kill microorganisms of root canal.^[3]

NaOCl is used as a proper irrigator in endodontics therapy that is effective against many bacteria and can solve necrotic pulp tissues. Toxicity for periapical tissues, inability in eliminating smear layer, change in physical structure of canal dentin and unfavorable smell and taste are some of disadvantages of NaOCl.^[4] Considering that without a proper root canal irrigation, root treatment will not be successful and since the root canal disinfection at the best condition cannot eliminate all microorganisms, the use of an irrigator, which has an antibacterial effect, and can close dentin tubules and neutralize bacteria with this way, is needed.^[2]

Cedar with scientific name, *Ziziphus spina-christi*, and English name, Lote tree or Christ's Thom is a plant of jujube family with 10 meter height, which grows self-growingly in Saudi Arabia, North Africa and Iran (in Khuzestan, Fars, Hormozgan provinces and Khark Island). Cedar is an evergreen plant with prickly branches, oval and taper leaves. Its fruit is sweet, sour, spherical, and aromatic, and has two-centimeter diameter. Its leaves powder is known as cedar and used

for hair irrigation and reinforcement as well as hair loss prevention. Another use of its leaf is disinfecting the wound. As mentioned before, water-alcohol extract of its leaf prevents the growth of some fungi. [5] Researches have shown that cedar leaf extract can have bactericidal effect against Gram-positive bacteria such as *Bacillus subtilis*, *Streptococcus pyogenes*, and *E. faecalis* as well as Gram-negative bacteria such as *Acinetobacter baumannii*, *Haemophilus influenzae*, *Salmonella enteritidis*, and *Escherichia coli*. [6] Since that cedar is a native plant of Iran and its south provinces, and its different pharmaceutical and antibacterial effects have been proved, the aim of this study was to compare antimicrobial effect of cedar and sodium hypochlorite 2.5% on *E. faecalis* bacterium.

Materials & Methods

1- Cedar leaf collection and extract preparation: The leaves of *Z. spina-christi* used in this study were collected in July 2017 from Ahvaz University of Medical Sciences in Khuzestan province of Iran. The leaves were dried at the shadow, room temperature for 48 h and crushed into powder. First, 90 gr of dried leaves were soaked in 800 ml of methanol and another 90 gr were soaked in 800 ml of ethanol 70% and allowed to stand in shaker (Labnet-America) for 72 h. Then, the solutions were filtered with No.42 What man filter paper. After that, solvents evaporated in oven (Shimi Fann-Iran) with 46°C for about 72 h. Finally, precipitates were scratched into powder using sterilized scalpel. [7]

2- 2.5% sodium hypochlorite: Solution of 5.25% sodium hypochlorite (chloraxid 5.25%- cerkamed) was diluted 1:2 with physiologic serum.

3-E. faecalis strain preparation: Standard strain of *E. faecalis* bacterium (ATCC 29212) was obtained from Iranian Research Organization for Science and Technology (Tehran-Iran).

4- Disk diffusion test: To evaluate antibacterial activity of *Z. spina-christi* leaves methanolic and hydro-alcoholic extracts and NaOCl, disk diffusion test was performed. First, standard strain of *E. faecalis* was inoculated in Tryptic soy broth (TSB) (Merck, Darmstadt, Germany) medium and incubated in 37 °C for 22 hours. Then, according to McFarland turbidometry, a suspension with 10⁸ CFU/ml final cell concentration was prepared from this culture. This microbial suspension was cultured on Mueller Hinton

agar plate (Merck, Darmstadt, Germany). Seven 6-mm diameter sterile paper disks (Padtanteb-Iran) were put on Mueller Hinton agar plate, and immediately 40 µl of each prepared concentration of methanolic or hydro-alcoholic extracts (0.05, 0.15, 0.25, 0.35, 0.45 g/ml), 20 µl of 2.5% NaOCl and 20 µl of physiologic serum (as negative control) were infused on two other paper disks. Plates were put in laboratory temperature for one hour till extracts were being absorbed to disks, then plates were incubated for 48 hours in 37 °C. [8] The inhibition zone around each disk was measured. This test was repeated three times and performed based on CLSI 2014 standard.

5- Microdilution test: To determine MIC (Minimum Inhibitory Concentration) and MBC (Minimum Bactericidal Concentration), culture of *E. faecalis* bacterium was diluted in sterile physiologic serum that a 10⁶ CFU/ml inoculum was gained with reference to 0.5 Mc Farland. In each well of microplate, serial dilution of methanolic and hydro-alcoholic extracts, 2.5% NaOCl and physiologic serum in 1:2 proportion was performed in Brain Heart Infusion (BHI) culture medium (Merck, Darmstadt, Germany) and 10 µl of standard suspension of *E. faecalis* was added to each well. After that, micro-plates were incubated in 37 °C. After 24 h, because of extract turbidity, MIC determining was not possible; therefore, we cultured 10 µl of each well at blood agar plates (Merck, Darmstadt, Germany) and incubated them in 37°C for 24 h. The lowest concentration of extract with less and no colony growth was considered as the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC), respectively. [9] This test was performed based on CLSI 2014 standard.

6- Statistical analyses: Data were analyzed using ANOVA. In case of significance, Scheffe's multiple comparison was used for two-by-two groups' comparison. P<0.05 was statistically considered significant.

Results

The results showed that hydro-alcoholic and methanolic extracts have antibacterial effect on *E. faecalis* bacterium and have better activity than physiologic serum as negative control (inhibition zone=0mm). The results of disc diffusion test indicated that inhibitory action of different concentrations of hydro-alcoholic and methanolic extracts against *E.*

faecalis bacterium was increased when used in higher concentration (figure-1).




Figure1. (a) Inhibition zone of methanolic extract and (b) Inhibition zone of hydro-alcoholic extract

Hydro-alcoholic extract has intermediate antibacterial activity on *E. faecalis* and it was more effective than methanolic extract (table-1). The inhibition zone of 2.5% NaOCl was significantly higher than other extracts ($p < 0.001$). Only one concentration (0.35g/ml) of methanolic extract was statistically comparable to hydro-alcoholic extract and 2.5% NaOCl. Mean inhibition zone of methanolic extract (0.35g/ml) was not significant ($p < 0.174$) compared to hydro-alcoholic extract (0.35g/ml), but it was significant in comparison with 2.5%NaOCl ($p < 0.001$). Mean inhibition zone of other concentrations (0.05, 0.15, 0.25

and 0.45 g/ml) of methanolic extract was 0, 0, 8 and 9 mm, respectively. The results of microdilution test demonstrated that *E. faecalis* bacterium was sensitive to both hydro-alcoholic and methanolic extracts but it was more sensitive to 2.5% NaOCl (table-2).

Table1. The comparison of mean inhibition zone (mm)

	Mean	SD	P-value*
NaOCl 2.5%	16.5	1.871	
Methanolic (0.05)	0	0.000	-
(0.15)	0	0.000	-
(0.25)	8	0.000	$P < 0.001$
(0.35)	8.33	0.577	$P < 0.001$
(0.45)	9	0.000	$P < 0.001$
Hydro-alcoholic (0.05)	8.33	0.577	$P < 0.001$
(0.15)	10	0.000	$P < 0.001$
(0.25)	10.67	0.577	$P < 0.001$
(0.35)	10.67	0.577	$P < 0.001$
(0.45)	11	1.000	$P < 0.001$

Table2. Minimum inhibitory and bactericidal concentrations of cedar methanolic and hydro-alcoholic extracts and 2.5% NaOCl on *E. faecalis*

	MIC	MBC
Cedar methanolic extract	62.5mg/ml	62.5mg/ml
Cedar hydro-alcoholic extract	62.5mg/ml	62.5mg/ml
2.5% NaOCl	0.625%	0.625%

Discussion

This study showed that methanolic and hydro-alcoholic extracts of *Z. Spina-christi* have antibacterial activity against *E. faecalis* in in vitro. Ramezanali et al.'s study in 2016 represented that green tea has acceptable antibacterial effect on the biofilm of *E. faecalis* [10], which is consistent with the study of Hudson et al. who found that cedar leaf (*Thuja pelicata*) oil, can kill *E. faecalis*. [6] The hydro-alcoholic extract than methanolic extract showed better inhibitory action, meaning it could be used more widely. According to Motamedi et al. in 2009, the ethanolic extract of *Z. spina-christi* illustrated slightly better killing action than methanolic extract against some Gram-positive and-negative bacteria, which are compatible with our results. [8] However, these both extracts showed lower potency compared to 2.5% NaOCl in both disc diffusion and microdilution tests. To

the best our knowledge, there is no study to comparatively evaluate the antibacterial activity of *Z. spina-christi* and 2.5% NaOCl against *E. faecalis*. Another study stated that the mean inhibition zone in descending order was found as sodium hypochlorite > Propolis > AI > Triphala > *C. longa* = MC > ethanol. [11] Recently, scientists from divergent fields are investigating plants for their antimicrobial usefulness. [12] The components of plants with antibacterial activity include tannins, saponins, phenolic compounds, essential oils and flavonoids. [13]

Bacteria are the most important factor in creating pulp and periapical disease. One of the bacteria in constant periapical lesions is *E. faecalis*. [1] The standard irrigator (2.5% NaOCl) used as positive control compared to the extracts showed higher antibacterial activity on *E. faecalis*. This is not surprising because it is well refined industrial product so there is no doubt its activity will be more than crude herbal extracts. If the extracts used in the present work were refined, more and better activity could be observed. [14] The major advantages of using herbal extracts alternatives are easy availability, low toxicity, increased shelf life and cost effectiveness. [15] This was a preliminary study that evaluated antibacterial activity of *Z. spina-christi* extract against *E. faecalis* in vitro; thus, further animal experimental studies are needed to determine its safety, toxicity, biocompatibility and efficacy. Considering disadvantages of 2.5% NaOCl which are mentioned previously and physiological benefits of this herbal extract, further studies using improved extraction technique and comparing common irrigants on other bacteria species can be carried out to consider it as an alternative for NaOCl at least in clinical situations like an open apex or in young permanent teeth, in which NaOCl causes toxicity.

Conclusion

As a result, 2.5% NaOCl had the highest antibacterial effect on *E. faecalis* followed by hydro-alcoholic and methanolic extracts. Therefore, NaOCl is an effective irrigator in root treatment until the studies like this can find a good alternative for it.

Acknowledgments

The authors would like to thank the Dental Materials Research Center for their supports.

Funding: This study was a part of research project (Grant no: 954318) which was supported and funded by Babol University of Medical Sciences. There is no conflict of interests.

Conflict of interest: The authors state that they have no conflict of interest.

Authors' contributions

The study was designed by Akam Saeidi and Mahmoodreza Hamidi. The study data was collected by Zahra Nikouee Rad. Analysis and interpretation of data drafting of the manuscript, and critical revision of the manuscript for important intellectual content were performed by Soraya Khafri, Akam Saeidi, Mahmoodreza Hamidi, Abolfazl davoodabadi and elham mahmoodi. Study supervision was conducted by Akam Saeidi and Mahmoodreza Hamidi and Abolfazl Davood abadi

References

1. Ehsani M, Amin Marashi M, Zabihi E, Issazadeh M1, Khafri S. A Comparison between Antibacterial Activity of Propolis and Aloe vera on *Enterococcus faecalis* (an In Vitro Study). *Int J Mol Cell Med* 2013; 2:110-6.
2. Razavian H, Mosleh H, Ehteshami A, Ziaei S, Nazeri R, Heidari F. Comparison of anti-bacterial effect of propolis and some conventional root canal irrigants. *J Isfahan Dent Sch* 2014; 10: 241-50. [In Persian]
3. Zare Jahromi M, kheibari M, rezaie S. [Effectiveness of sodium hypochlorite and chlorhexidine irrigation in prevention of longterm root canal periapical seal microlicage]. *J Res Dent Sci* 2009; 6: 24-7. [In Persian]
4. Jafari A, Kiani M, Nikkhah M, Bakhtiari R, Baradaran Nakhjavani Y, Heidari A R. Comparative evaluation of antiseptic effects of sodium hypochlorite, thyme essence and normal salin in root canal irrigation of primary teeth. *J Dent Med-Tehran Univ Med Sci* 2014; 27:161-7. [In Persian]
5. Gharib Naseri M K. Effect of hydroacoholic *Zizyphus spina christi* leaf extract on perfused frog heart contractility and frequency. *Feyz* 2003; 7: 1-7. [In Persian]
6. Hudson J, Kuo M, Vimalanathan S. The antimicrobial properties of cedar leaf (*thuja plicata*) oil; a safe and

- efficient decontamination agent for buildings. *Int J Environ Res Public Health* 2011; 8: 4477-87.
7. Mirpour M, Gholizadeh Siahmazgi Z, Sharifi Kiasaraie M. Antibacterial activity of clove, gall nut methanolic and ethanolic extracts on *Streptococcus mutans* PTCC 1683 and *Streptococcus salivarius* PTCC 1448. *J Oral Biol Craniofac Res* 2015; 5: 7–10.
 8. Motamedi H, Safary A, Maleki S, Seyyednejad S M. *Ziziphus spina-christi*, a native plant from khuzestan, iran, as a poteneial source for discovery new antibacterial agents. *Asian J Plant Sci* 2009; 8: 187-90.
 9. Ebadi AR, Monadi A, Pashazadeh M, Zakhireh S. Antibacterial effects of *Astragalus hamosus* ethanolic extract on some gram positive and negative bacteria. *J Comp Pathol Iran* 2014; 10: 1071-6.
 10. Ramezanali F, Samimi S, Kharazifard M , Afkhami F. The in vitro antibacterial efficacy of persian green tea extract as an intracanal irrigant on enterococcus faecalis biofilm. *Iran Endod J* 2016;11: 304-8.
 11. Saxena D, Saha SG, Saha MK, Dubey S, Khatri M. An in vitro evaluation of antimicrobial activity of five herbal extracts and comparison of their activity with 2.5% sodium hypochlorite against *Enterococcus faecalis*. *Indian J Dent Res* 2015; 26:524-7.
 12. Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev* 1999; 12: 564-82.
 13. Karkare SR, Ahire NP, Khedkar SU. Comparative evaluation of antimicrobial activity of hydroalcoholic extract of Aloe vera, garlic, and 5% sodium hypochlorite as root canal irrigants against *Enterococcus faecalis*: An in vitro study. *J Indian Soc Pedod Prev Dent* 2015; 33:274-8.
 14. Abalaka M E, Daniyan S Y, Mann A. Evaluation of the antimicrobial activities of two ziziphus species (*ziziphus mauritiana* l. and *ziziphus spinachristi* l.) on some microbial pathogens. *Afr J Pharm Pharmacol* 2010; 4: 135-39.
 15. Chaitanya BV, Somisetty KV, Diwan A, Pasha S, Shetty N, Reddy Y, et al. Comparison of antibacterial efficacy of turmeric extract, morinda citrifolia and 3% sodium Hypochlorite on *Enterococcus faecalis*: An In-vitro Study. *J Clin Diagn Res* 2016; 10: ZC55-7.