

e-ISSN: 2322-2395 p-ISSN: 2251-9890



Knowledge and therapeutic approach of Iranian dentists toward COVID-19 pandemic Nika Mehrnia^{1⊠}, Reyhaneh Eghbali Zarch^{1⊠}, Maryam Sadeghipour^{2*}

1. Postgraduate Resident, Department of Periodontics, Dentistry Faculty, Shahed University, Tehran, Iran.

2. Assistant Professor, Department of Community Oral Health, School of Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Article Type	ABSTRACT			
Research Paper	Introduction: Dental services are a hotspot for viral transmission. Dentists must have complete			
	knowledge about the disease and quick access to information to combat this transmission. To prevent			
	COVID-19 viral spreading through dental practices, dentists must be well-versed in disease and infection			
	control measures. As a result, this study aimed to assess the knowledge of Iranian dentists on the Covid			
	19 pandemic.			
	Materials & Methods: The present study was a questionnaire-based cross-sectional survey. A modified			
	questionnaire of a perused one was prepared, the validity of which was confirmed, and the reliability was			
	accepted by Cronbach's alpha test (0.86). This questionnaire was completed by 133 Iranian dentists and			
	included six components: consent to participate in research, demographic information, knowledge,			
	infection control, vaccination, and therapeutic approach. Data were analyzed using Mann-Whitney,			
	Kruskal-Wallis, and Spearman rank correlation coefficients, and P<0.05 was considered the level of			
	significance.			
	Results: Iranian dentists are shown to have proper knowledge, but only 21.7% of participants received			
	infection control courses; moreover, only 47% of them knew the accurate incubation period of the Virus.			
	Dentists working in Academic clinics took more preventive measures than those working in private clinics			
	(P=0.023).			
Received: 13 Oct 2021	Conclusion: It is recommended that health care authorities ensure that dentists receive organized			
Revised: 13 Nov 2021	information and guidelines.			
Accepted: 16 Mar 2022	Keywords: COVID-19, Dentists, Infection Control, Knowledge, Surveys and Questionnaires			
Cite this article: Mehrnia N, Eghbali Zarch R Sadeghipour M. Knowledge and therapeutic approach of Iranian dentists toward				
COVID-19 nandemic	Caspian I Dent Res 2022: 11(1):46-55			

© The Author(s). Publisher: Babol University of Medical Sciences

Introduction

Coronavirus is a term that refers to a range of respiratory viruses that can cause mild to severe symptoms and lead to respiratory failure. Its name recalls the virus's morphology, including pointed structures on the surface that resemble

E-mail: sadeghipour1393@gmail.com

^{*} Corresponding Author: Maryam Sadeghipour, School of Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Tel: +982122175350

a crown.^[1]The origin of SARS-CoV-2 is unknown, although it is believed that the virus was first transmitted from animals to people in the Huanan seafood market in Wuhan, China, in December 2019, where a cluster of pneumonia cases caused by a newly found coronavirus was diagnosed.^[2] It was declared a pandemic emergency by World Health Organization (WHO) in March 2020. Common clinical features of this virus infection include fever, cough, headache, diarrhea, fatigue, headache, and myalgia. Anosmia and ageusia are other commonly reported symptoms.^[3]

COVID-19 has an incubation time that ranges between 1 and 14 days. Early detection of symptoms is life-saving. Some mucocutaneous manifestations of this viral infection can be beneficial for the early diagnosis. There were also occasions where the only manifestation was oral symptoms, and some reports indicated that approximately 80% of infected cases were asymptomatic.^[4, 5] Many regular dental treatments can transfer the virus through aerosols, making dentists one of the most significant risk groups for transmission and infection with the coronavirus. As a result, the demand for personal protective equipment has increased. In case of airborne infections like COVID-19, personal protective equipment consists of gloves, masks, gowns, face protection, head cover, and rubber boots.^[6]

Iran was one of the first countries infected with the virus after the epidemic in Wuhan, China.^[7] The first cases in Iran were reported on 19 February 2020 in Qom.^[8] Soon after that, many more cases were identified in other provinces, with some provinces being highly affected in the early phase of the epidemic, including Tehran, Gilan, Mazandaran, Qom, and Golestan.^[9] Due to the latest report by the public relation and information center of the Ministry of Health of Iran in November 2021, a total of 6 million COVID-19 positive cases were reported in Iran, about 128000 of whom have died. Nevertheless, more than 3000 COVID-19 positive patients were being cared for in hospital intensive care units at the time of the report. ^[10] To combat the outbreak of COVID-19, dentists must have a high degree of knowledge about the disease, its severity, and infection control measures. Various studies were conducted in different countries to assess their dentist's knowledge. In this regard, this study was conducted to assess Iranian dentists' knowledge, attitudes, and prevention strategies regarding the COVID-19 pandemic.

Materials & Methods

The present study is a questionnaire-based cross-sectional survey approved by code IR.SBMU.DRC.REC.1399.151 in the Research Council of Shahid Beheshti University of Medical Sciences. Due to COVID-19 pandemic circumstances, the actual number of active dentists was unavailable, making a precise assessment of the target population number impossible. As a result, from May 22nd to June 27th, 2020, this study was conducted using Google Forms and circulated through social media groups.

Sample size: The statistical population was the target of dentists in the whole country, the total number of which was announced by the Ministry of Health as nearly 20,000 dentists. Consequently, the statistical population had a limited statistical opinion and many changes. The following formula is used to determine the sample size:

$$n = \frac{N * Z_{\frac{\alpha}{2}}^2 * \sigma^2}{\varepsilon^2 (N-1) + Z_{\frac{\alpha}{2}}^2 * \sigma^2}$$

Where N represents the volume of the target population and is equal to 20,000, α is the level of error and equal to 0.05, ϵ is the accuracy of the researcher, which usually takes a value equal to 10%, and σ is the standard deviation equal to 0.59.^[11] As a result, the number of samples required for this study was obtained to be 132.84 people, which was considered by rounding 133 people. It should be noted that the number of active dentists reported by the Ministry of Health (20,000) is based on pre coronavirus conditions. In addition, many dentists did not work because of the corona pandemic; therefore, the exact number of active dentists during the pandemic was not known.

Finally, 133 dentists participated in the study. Participants provided consent regarding publishing their data in the first section of the questionnaire, their personal information was protected, and they had the option to discontinue responding to the survey at any time. The questionnaire was a modified version of the one used by Mustafa et al.^[12]. Two different experts evaluated the original questionnaire, selected appropriate questions, and modified them. The validity of this questionnaire was reviewed through the content validity method by six professors in this field and

approved after making the desired corrections. Moreover, Cronbach's alpha was used to evaluate the reliability of the questionnaire, and it was 0.86, which is an acceptable value, confirming the strong reliability of this questionnaire.

The questionnaire contained 32 questions in 5 sections, including demographic information, knowledge, dental treatments, vaccine, and risk assessment. SPSS version 26 was used to analyze the data. The relationship between sex and other categorical variables was assessed using the Mann-Whitney test. Also, the Kruskal-Wallis test was employed to evaluate the relationship between age, practice location, degree of education, and other categorical variables. Other than demographic data, the Spearman rank correlation coefficient was used to explore any correlation between different questionnaire parts. The statistical significance level was set as a P value of less than 0.05.

In order to evaluate the questionnaire, every correct answer was considered as 1 point. However, some questions had more than one correct answer, which could result in more than 1 point; false answers did not end with a negative point. Finally, the sums of points were calculated in each section, the average of which was used to evaluate and compare study groups. The overall points in each section were as follows: Knowledge: 14, Dental treatments and infection control: 12, Vaccine: 5, Risk assessment and therapeutic approach: 10.

Results

Demographic Data: A total of 133 responses were collected from Iranian dental practitioners. The majority of responders were female, with an average age of 20-29. Most of the participants were general dentists who worked in private practices. Table 1 presents the demographic characteristics of the participants in detail.

Table 1. Demographic data of participant	5
Demographic Variables	percent
Gender	
Male	26.6
Female	73.4
Age	
20-29	71.1
30-39	21.9
40-49	3.1
50-59	3.9
\geq 1.	0
Region of practice in the country	
Central region	42.1
Northern region	27
Western region	18.3
Eastern region	4
Southern region	8.7
Working sector	
University and dental schools	35.7
Private practice	46.5
University clinics	17.1
Others	0.8
Education Status	
Dentistry student	20.9
General dentist	60.5
Specialist/Resident	18.6
Participation in COVID-19 infection control courses	
Yes	21.7
No	78.3

Caspian Journal of Dental Research, March 2022; 11(1): 46-55

Knowledge: The participants demonstrated proper knowledge of COVID-19 symptoms with an average point of 8.47. Also, 91.7% of dentists agreed that fever and cough are the most common disease symptoms. Moreover, 96.9% were aware that the virus was spread by aerosol, and 68.4% answered the contact transmission chance correctly. However, only 47.3% of participants knew the answer to the incubation period, and only 21.7% participated in infection control courses. Although dental residents and specialized dentists had a little more knowledge about COVID-19 and got 8.8 points compared to general dentists, who got 8.3 points on average, the difference was not statistically significant (P=0.802). The outcome of the knowledge section is presented in Table 2.

Table 2. Participant responses to the knowledge section	
Question	
Which one is the incubation period of the coronavirus?	
2-7 days	22.5
2-14 days	47.3
7-14 days	22.5
7-21 days	7.8
Which of the following is a COVID-19 symptom? (There might be multiple right answers.)	
Fever	96.2
Cough	91.7
Runny nose	41.3
Sore throat	66.9
Shortness of breath	52.6
Joint / muscle pain	38.3
Eye redness	15
Skin rash	20.3
Diarrhea	33.8
Vomit	26.3
It may be asymptomatic	39.8
Which of the following is a COVID-19 transfer method? (There might be multiple right answers.)	
Respiratory (through sneezing and coughing)	96.9
Handshaking	68.4
Touch surfaces (door handles, tables, etc.)	47.3
Which of the following is considered a risk for COVID-19? (There might be multiple right answers.)	
Diarrhea symptoms	50.3
Respiratory infection symptoms	84.9
History of travel to high prevalence areas or new species of COVID-19	85.7
History of Contact with the infected person	93.2

Dental Treatments and Infection Control: The majority of responders used appropriate personal protective equipment with an average point of 5.9. In addition, 96% believed that personal protective equipment is effective against viral transmission. Face protection was beneficial to 97.7% of dentists, and 85.3% used a combination of two masks; however, most dentists did not use COVID-19 rapid test, pulse oximeter, and thermometer before treatment. Table 3 presents the dental treatments and infection control measures section.

Vaccine: Most of the participants had received the COVID-19 vaccination and were well versed in the subject, gaining an average of 3.5 points. 82% of individuals had received their vaccination at the study time. Moreover, dentists with higher knowledge of vaccination were more cautious in their infection control measures and treatment approaches (P<0.05). Although dental residents and specialist dentists had higher knowledge about the COVID-19 vaccine (3.7 points), the difference was not statistically significant compared to general dentists, who gained 3.3 points on average (P=0.128). (Supplementary Table 1)

Table 5. Dentists Tesponses regarding infection control measures	
Question	
The use of personal protective equipment such as shields and goggles, FFP2 masks (KN95) and	
gloves is effective in protecting medical staff in the face of a patient suspected of COVID-19.	
Yes	96.9
No	0
No idea	3.1
How many masks do you use during your clinical activity?	
A three-layer surgical mask	4.7
Two three-layer surgical masks	27.9
An N95 mask and a three-layer surgical mask	57.4
3M mask	7.8
Others	2.3
How much is the risk of transmitting the infection reduced by using a cloth mask on a three-layer	
mask or fitting the mask by tying it?	
60 percent	16.8
70 percent	28
80 percent	16.8
95 percent	38.4
Which of the following protective equipment can you use during dental treatment?	
Goggles	16.2
Simple face shield	50
Casket hat face shield	31.5
I do not use these devices	2.3
Do you get Rapid Tests from your patients before starting treatment?	
Yes	1.6
No	98.4
Do you use a pulse oximeter for your patients before starting treatment?	
Yes	21.1
No	78.9
Do you measure your patients' body temperature before starting treatment?	
Yes	38.3
No	61.7
Do you use ultrasonic devices for SRP during the Pandemic period?	
Yes	54.4
No	45.6

Question	
Have you received the vaccination?	
Yes	82
No	28
How long does it take immunity to be developed after vaccination is given?	
Immediately after the first dose injection	2.3
Immediately after the second dose injection	
Sometime after the first dose injection	27.1
Sometime after the second dose injection	69.8
If the vaccination is fully effective, which of the following might be possible? (There might be	
multiple right answers.)	
Severe form of the disease or being hospitalized due to COVID-19	9.7
Mild type infection	
Vaccinated person may be a carrier of a virus	56.3
what percentage of the population should be vaccinated in order to break the transmission chain?	
100%	7.8
80%	59.7
60%	31
30%	1.6
Is vaccination effective against the new species?	
Yes	76.6
No	23.4

Risk assessment and Therapeutic Approach: Based on the comments of participants, COVID-19 is a critical health concern that requires special consideration while treating dental patients. There is a significant relationship between dentists' place of practice and their therapeutic approach to COVID-19. Dentists working in universities clinics took significantly more preventive measures in their therapeutic approaches with an average of 4.4 compared to those who worked in their private practice with an average of 3.8 (P value<0.05). Furthermore, females were shown to be more conservative in their therapeutic approach, gaining an average of 4.11 compared to men, who gained 3.79; however, the difference was not statistically significant (P value=0.150). Supplementary Table 2 presents the risk assessment section. It is worth noting that there was no evidence of a significant relationship between other category variables.

Supplementary Table 2. Responses of participants to the risk assessment and therapeutic approach section

Question	
How do you assess the risk of COVID-19?	
High risk	67.4
Medium risk	32.6
Mild risk	0
Do you consider COVID-19 an important health issue?	
Yes	98.4
No	1.6
The symptoms of COVID-19 go away over time and there is no need for special treatment.	
Yes	21.7
No	78.3
What do you do if a patient comes to you with symptoms of coughing or sneezing?	
Failure to perform treatment and asking the patient to leave the center	35.7
Performing treatment and requesting the patient to go to the hospital	3.9

Referral of the patient to the hospital without treatment	60.5
What is your opinion about maintaining distance between patients and the need for them to use the	
mask while waiting for a visit and hand washing before entering the center?	
It is essential and helps reduce the spread of the disease	100
It is not necessary but it may cause fear	0
If one of your clinic staff members has flu-like symptoms, will you allow them to work with the	
patient?	
Yes	1.6
No	98.4
What do you think is the role of dentists in educating others about COVID-19?	
Very high	35.7
Medium	54.3
Mild	9.3
None	0.8
If you are in contact with an infected person, do you know where to go or who to contact?	
Yes	77.3
No	22.7
If you have any suspicious symptoms of COVID-19, do you know where to go or who to contact?	
Yes	82.5
No	17.5

Discussion

Due to the proximity of dentists to the working field, they are highly susceptible to the COVID-19 virus. The present study revealed that Iranian dentists have proper general knowledge about COVID-19; however, only 21.7% of participants received infection control courses. Moreover, only 47% of them knew the accurate incubation period of the virus, which is in accordance with some studies conducted in other countries.^[12, 13] While showing proper general knowledge may reflect the high motivation of dentists to learn about the issue, a lack of information in a specific area may indicate that the correct information was not easily accessible. On the other hand, Sarfaraz et al. conducted a multi-country study in four countries, revealing insufficient knowledge in fundamental aspects of COVID-19 among dentists while having a positive attitude toward disinfection practices during this pandemic. To summarize the results of the studies mentioned above, it is strongly recommended that health care authorities provide organized materials that dentists can easily access, especially since we had been dealing with the pandemic for more than a year at the time of the study.^[14]

Over 95% of participants used proper infection control equipment, which is critical in preventing COVID infection. According to the research findings, there is a significant difference in the therapeutic approaches among dentists working in different areas. Dentists working in universities clinics had more preventive measures in their therapeutic approaches, attributed to dental school administrators' attitudes toward meticulous patient screening to avoid delivering dental treatments to COVID-19 patients. In order to minimize the risk of disease transmission through dental practices, it seems this restrictive attitude should be extended to all dentists across the country, which can be accomplished by motivating them to follow CDC and WHO guidelines. Several pieces of previous studies have shown that better knowledge was associated with higher prevention measures.^[15-17] This association was not significant in the present study, and dentists who practice in university clinics have shown much more preventive therapeutic approaches. Naturally, some discrepancies in the results of different studies might be related to the time they were conducted; some studies were performed early in the pandemic with limited information or guidelines.^[12]

In the present study, postgraduate residents and dentists had slightly higher knowledge scores than graduates and undergraduates. This finding is in line with other worldwide surveys ^[16, 18-21] and pieces of research regarding previous pandemics such as ZIKV and Ebola.^[22, 23] Involvement of graduates in research and exploration of the latest pieces of

evidence could be the reason for this. On the other hand, Khanal et al. found no significant association between education levels and COVID-19 related knowledge scores, and Harapan et al. reported higher scores during the Zika virus pandemic for graduates compared to higher education.^[24, 25] The disparity in the findings of these two studies could be attributed to differences in education curriculum and rules between countries.

Ahmed et al. reported anxiety and fear among dental practitioners around the globe; participants were afraid of getting infected with COVID-19 from either a patient or a co-worker or of spreading the virus from dental practice to their families. There was no statistical relationship between the responses and gender in their study. ^[26] In contrast, in the present study, women were found to be slightly more conservative when providing dental treatments to patients with COVID-19 like symptoms. Anxiety and obsessive-compulsive symptoms are psychological dangers for healthcare personnel during the pandemic.^[27] These findings could be consistent with those of other studies that have suggested higher levels of job-related anxiety in women.^[28, 29]

Leadership scholars have widely recommended learning from experiences. ^[30, 31] Countries that used their prior experience dealing with emerging infections, such as China and South Korea, have more efficiently responded to the COVID-19 pandemic. Therefore, the results of this study can be helpful during the rest of the COVID-19 era and can be used as a learning tool from experience in future pandemics.

The limitation of the present study was that participants were contacted via social media, which was unavoidable since COVID-19 prohibited all scientific gatherings. Nonetheless, the cross-sectional design of this study precludes making causal inferences.

Conclusion

This cross-sectional study revealed that Iranian dentists have the proper knowledge and therapeutic approaches regarding COVID-19; however, there is a lack of knowledge on the disease incubation period. Only a small number of participants got infection control courses during the pandemic. As a result, to limit infection transmission through dental practices, healthcare authorities are suggested to ensure that organized information and guidelines are provided to dentists. Furthermore, the findings of this study could provide strategies for healthcare authorities to take faster measurements and be better prepared for future epidemics and pandemics.

Conflicts of Interest

There is no conflict of interest

Authors' Contribution

N. M. contributed to the conception, design, data acquisition, interpretation, statistical analysis as well as drafted and critically revised the manuscript.

R. E. Z. contributed to the conception, design, data acquisition, interpretation, statistical analysis, as well as drafted and critically revised the manuscript.

M. S. contributed to the conception, design, data acquisition, interpretation, statistical analysis as well as drafted and critically revised the manuscript.

All of the authors have read and approved the submitted version of the manuscript.

References

Yang Y, Peng F, Wang R, Yange M, Guan K, Jiang T, et al. The deadly coronaviruses: The 2003 SARS pandemic and the 2020 novel coronavirus epidemic in China. J Autoimmun 2020;109:102434.
Brown JD. Antihypertensive drugs and risk of COVID-19 ? Lancet Respir Med2020;8:e28.

Caspian Journal of Dental Research, March 2022; 11(1): 46-55

3. Kara C, Çelen K, Dede FÖ, Gökmenoğlu C, Kara NB. Is periodontal disease a risk factor for developing severe COVID-19 infection? The potential role of Galectin-3. Exp Biol Med (Maywood) 2020;245:1425-7.

4. Day M. COVID-19 : four fifths of cases are asymptomatic, China figures indicate. BMJ 2020; 369:m1375.

5. Eghbali Zarch R, Hosseinzadeh P. COVID-19 from the perspective of dentists: A case report and brief review of more than 170 cases. Dermatol Ther2021;34:e14717.

6. World Health Organization. Rational use of personal protective equipment for coronavirus disease (COVID-19) and considerations during severe shortages. Available at: file:///C:/Users/asri/Downloads/WHO-2019-nCoV-IPC_PPE_use-2020.4-eng.pdf. Accessed June 2020.

7. Mounesan, L, Eybpoosh S, Haghdoost AA, Moradi, G, Mostafavi E. Is reporting many cases of COVID-19 in Iran due to strength or weakness of Iran's health system? Iran J Microbiol 2020; 12: 73.

8.Doosti-Irani A, Haghdoost AA, Najaf F, Eybpoosh S, Moradi G, Bagheri Amiri F. How can the epidemic curve of COVID-19 in Iran be interpreted? J Res Health Sci2020; 20: e00491.

9. Moradi Y, Eshrati B. Estimation of the net reproductive number of COVID-19 in Iran. Med J Islam Repub Iran 2020; 34:34.

10. Ministry of Health and Medical Education. Statistics of patients with COVID-19 . 2021[Online]. Aviailable From: https://behdasht.gov.ir

11.Miot HA. Sample size in clinical and experimental trials. J Vasc Bras 2011; 10: 275-8.

12.Mustafa RM, Alshali RZ, Bukhary DM. Dentists' Knowledge, Attitudes, and Awareness of Infection Control Measures during COVID-19 Outbreak: A Cross-Sectional Study in Saudi Arabia. Int J Environ Res Public Health 2020; 17: 9016.

13.Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK. Knowledge and Perceptions of COVID-19 Among Health Care Workers: Cross-Sectional Study. JMIR Public Health Surveill2020;6:e19160.

14. Sarfaraz S, Shabbir J, MudasserMA, Khurshid Z, Al-Quraini AAA, Abbasi, MS,et al. Knowledge and attitude of dental practitioners related to disinfection during the COVID-19 pandemic. Healthcare 2020;8:1-9.

15.Hleyhel M, Haddad C, Haidar N, Charbachy M, Saleh S. Determinants of knowledge and prevention measures towards COVID-19 pandemic among Lebanese dentists: a cross sectional survey. BMC Oral Health 2021; 21: 241.

16. Kamate SK, Sharma S, Thakar S, Srivastava D, Sengupta K, Hadi AJ, et al. Assessing Knowledge, Attitudes and Practices of dental practitioners regarding the COVID-19 pandemic: A multinational study. Dent Med Probl 2020;57:11-7.

17. Abou-Abbas L, Nasser Z, Fares Y, Chahrour M, El Haidari R, Atoui R. Knowledge and practice of physicians during COVID-19 pandemic: a cross-sectional study in Lebanon. BMC Public Health 2020;20: 1474.

18. Ramesh A, Nagalaxmi V, Patel BP. Knowledge and Awareness of Oral Health Care Professionals Regarding Practices and Controls in Dental Practice Following the COVID-19 Outbreak in the State of Telangana-A Questionnaire Study. Int J Innov Sci Res Technol

2021;6:432-8.

19. Chowdhury MTH, Hoque Apu E, Nath SK, Noor AE, Podder CP, Mahmud I,et al.

Exploring the knowledge, awareness and Practices of COVID-19 among dentists in Bangladesh: A cross-sectional investigation. J Oral Res 2020;10:1-12.

20.Singh Gambhir R, Singh Dhaliwal J, Aggarwal A, Anand S, Anand V, Kaur Bhangu A.COVID-19 : a survey on knowledge, awareness and hygiene practices among dental health professionals in an Indian scenario. Rocz Panstw Zakl Hig 2020; 71:223-9.

21.Quadri MFA, Jafer MA, Alqahtani AS, Al Mutahar SAB, Odabi NI, Daghriri AA, et al. Novel corona virus disease (COVID-19) awareness among the dental interns, dental auxiliaries and dental specialists in Saudi Arabia: A nationwide study J Infect Public Health 2020;13:856-64.

22.Gupta N, Randhawa RK, Thakar S, Bansal M, Gupta P, Arora V. Knowledge regarding Zika virus infection among dental practitioners of tricity area (Chandigarh, Panchkula and Mohali), India. Niger Postgrad Med J 2016;23:33-7.

23. Holakouie-Naieni K, Ahmadvand A, Raza O, Assan A, Elduma AH, Jammeh A, et al. Assessing the Knowledge, Attitudes, and Practices of Students Regarding Ebola Virus Disease Outbreak. Iran J Public Health 2015;44:1670.

24. Khanal N, Singh AK. Knowledge, Attitude and Practice Regarding COVID-19 and its Impact on Dentistry: A Cross-sectional Survey among Nepalese Dentists. Kathmandu Univ Med J (KUMJ) 2020;18:3-9.

25. Harapan H, Aletta A, Anwar S, Setiawan AM, Maulana R, Wahyuniati N, et al. Healthcare workers' knowledge towards Zika virus infection in Indonesia: A survey in Aceh. Asian Pac J Trop Med 2017;10:189-94.

26. Ahmed MA, Jouhar R, Ahmed N, Adnan S, Aftab M, Zafar MS, et al. Fear and practice modifications among dentists to combat novel coronavirus disease (COVID-19) outbreak. Int J Environ Res Public Health2020; 17: 2821.

27. Goulia P, Mantas C, Dimitroula D, Mantis D, Hyphantis T. General hospital staff worries, perceived sufficiency of information and associated psychological distress during the A/H1N1 influenza pandemic. BMC Infect Dis 2010;10:322.

28. Choy HB, Wong MC. Occupational stress and burnout among Hong Kong dentists. Hong Kong Med J 2017; 23: 480-8.

29. Tysiąc-Miśta M, Dziedzic A.The attitudes and professional approaches of dental practitioners during the COVID-19 outbreak in Poland: a cross-sectional survey. Int J Environ Res Public Health 2020; 17: 4703.

30. Thomas RJ. Crucibles of leadership: How to learn from experience to become a great leader. Brighton, Massachusetts: Harvard Business Publishing,;2008.p.

31. The Lancet. COVID-19: learning from experience. Lancet 2020; 395:1011.