

## Evaluation of soft tissue norms on lateral cephalograms in babol

Valiollah Arash<sup>1</sup>, Manouchehr Rahmati Kamel<sup>2</sup>, Abbasali Ostad Rahimi<sup>3</sup>, Reza Ghorbanipour<sup>2✉</sup>

1. Associate Professor, Dental Materials Research Center, Department of Orthodontics, Faculty of Dentistry, Babol University of Medical Sciences, Babol, IR Iran.
2. Assistant Professor, Dental Materials Research Center, Department of Orthodontics, Faculty of Dentistry, Babol University of Medical Sciences, Babol, IR Iran.
3. General Dentist, Gorgan, IR Iran.

✉**Corresponding Author:** Reza Ghorbanipour, Faculty of Dentistry, Babol University of Medical Sciences, Babol, IR Iran.

**Email:** dr\_ghorbanipour@yahoo.com

**Tel:** +98 1132291408-9

**Received:** 2 May 2017      **Accepted:** 26 Sept 2017

### Abstract

**Introduction:** Soft tissue has a prominent role in diagnosis and treatment plan in orthodontics. Facial proportions are measurable on lateral cephalograms. The purpose of this study was to compare the means of the normal soft tissue proportions of babol people with the norms of Caucasians.

**Materials & Methods:** In this cross-sectional study, 100 cases with normal occlusion and proportional facial profile were participated from babol. After taking radiographs and tracing, the soft tissue variables were analyzed to find means and standard deviations. The data were compared with Caucasian's norms using T-test.

**Results:** The norms of following variables in babol people were significantly different from those of Caucasians: facial convexity, nasolabial angle, upper lip thickness, chin soft tissue thickness and protrusion of lower lip.

**Conclusion:** In babol, people have a more convex profile, more prominent nose and lower lip than caucasians.

**Keywords:** Cephalometry, Tissue, Treatment, Orthodontics

*Citation for article:* Arash Vo, Rahmati Kamel M, Ostad Rahimi Aa, Ghorbanipour R. Evaluation of soft tissue norms on lateral cephalograms in babol. Caspian J Dent Res 2017; 6: 30-4.

## ارزیابی مقادیر نرمال بافت نرم در رادیوگرافی های لترال سفالومتری در شهر بابل

ولی الله آرشی، منوچهر رحمتی کامل، عباسعلی استاد رحیمی، رضا قربانی پور\*

### چکیده

**مقدمه:** بافت نرم نقش مهمی در تشخیص و طرح درمان در ارتودنسی دارد. نسبت های صورتی از طریق رادیوگرافی لترال سفالومتری قابل اندازه گیری هستند. هدف از این مطالعه، مقایسه میانگین نسبت های بافت نرم نرمال در شهر بابل با مقادیر نرمال در کاکازینها می باشد.

**مواد و روش ها:** در این مطالعه مقطعی، ۱۰۰ نفر با اکلوزن و نیم رخ با نسبت های نرمال از شهر بابل شرکت کردند. بعد از گرفتن رادیوگرافی لترال سفالومتری و انجام تریسینگ، متغیرها جهت یافتن میانگینها و انحراف معیار آنالیز شدند. داده ها با مقادیر نرمال کاکازین به کمک T test مقایسه شدند.

**یافته ها:** اعداد به دست آمده برای متغیرهای زیر با مقادیر نرمال در پروفایل بافت نرم نرمال کاکازینها متفاوت بودند: تحدب صورت، زاویه نازولیبال، ضخامت لب بالا، ضخامت بافت نرم چانه و بیرون زدگی لب پایین.

**نتیجه گیری:** در بابل، افراد نرمال نیمرخ محدبتر و بینی و لب پایین برجسته تری نسبت به کاکازینها دارند.

**واژگان کلیدی:** سفالومتری، بافت، درمان، ارتودنسی


### Introduction

**E**valuation of soft tissue in patients demanding esthetic procedures such as orthodontics and orthosurgery has a prominent role in diagnosis and treatment planning. Beforehand, hard tissue and teeth were in focus in orthodontics and orthosurgery and now, soft tissue proportions have higher importance. <sup>[1, 2]</sup> According to the importance of facial proportions in esthetics and patient's expectation, the efforts have led to evaluate the soft tissue proportions. <sup>[3]</sup> At first, means of soft tissue proportions were driven from Caucasian population and considered as a criterion for comparison in cephalometric analysis. <sup>[1, 4- 6]</sup> Afterwards, several researches on different ethnics suggested the differences between Caucasian's norms and norms of other races. <sup>[7- 11]</sup> Racial studies evaluated the norms of soft tissue proportions in different Iranian populations. A study in Ahwaz showed a more convex profile and more prominent nose in Khuzestan compared to European standards. Lips and chin are more retruded than nose in this area. <sup>[12]</sup> Moghbel et al. in Tehran found that facial profile is more convex and chin soft tissue is thicker in Iranian population than Caucasians. <sup>[13]</sup> There is no published data on the soft tissue norms in babol. Therefore, the purpose of this study was to evaluate the


soft tissue proportions of normal facial profile in Babol to drive means in this region and to compare these normative data with norms of Caucasian race.

### Materials &Methods

One hundred of students and adult patients (34 males and 66 females) aged from 20 to 30 years, <sup>[8]</sup> referred to babol school of dentistry participated in this cross-sectional descriptive- analytical study. All persons were from mazandaran and had normal facial proportions in clinical evaluation, normal class I occlusion, minimal anterior crowding, without history of orthodontics and esthetic procedures and surgeries. After getting informed consent, all cases were referred to one radiologic center with a specific magnification by a single technician. Cephalograms were in neutral head position with lips in rest and teeth in maximum intercuspation. All the steps of tracing and measurement of variables were done by a single orthodontist. According to Holdaway analysis <sup>[14]</sup> (Fig. 1), Epker's soft tissue relations <sup>[15]</sup> (Fig. 2) and Legan-burstone soft tissue analysis <sup>[4, 5]</sup> (Fig. 3), soft tissue variables were measured.




**Fig 1. Holdaway soft tissue analysis measurements.** A, soft tissue facial angle; B, nose prominence; C, superior sulcus depth; D, soft tissue subnasale to H line; E, skeletal profile convexity; F, upper lip thickness; G, upper lip strain; H, H angle; I, lower lip to H line; J, inferior sulcus to H; K, soft tissue chin thickness



**Fig 2. Epker's soft tissue relations.** G, glabella; Sn, subnasale; St, stomion; ME, menton

All data were analyzed using the statistical package for social sciences (version 11; SPSS Inc., Chicago, Illinois, USA). Means and standard deviations were calculated for each measurement. To compare Mazandarani and Caucasian norms, independent T-tests were used.  $P < 0.05$  was considered as significant level.



**Fig 3. Legan-Burstone soft tissue analysis.** Horizontal reference plane (HP), constructed by drawing a line through nasion (N) 7 degrees up from the sella-nasion line. 1, facial convexity angle (G-Sn-Pg'); 2, maxillary prognathism (G vertical-Sn); 3, mandibular prognathism (G vertical-Pg'); 4, lower face-throat angle (Sn-Gn'-C); vertical height ratio (G-Sn/Sn-Me'), lower vertical height-depth ratio (Sn-Gn'/C-Gn'), nasolabial angle (Cm-Sn-Ls), upper lip protrusion (Ls to Sn-Pg'), lower lip protrusion (Li to Sn-Pg'), mentolabial sulcus (Si to Li-Pg'), vertical lip-chin ratio (Sn-Stms/Stmi-Me'), maxillary incisor exposure (Stms-UI), interlabial gap (Stms-Stmi).

## Results

Means and standard deviations of each variable in all cases of the study are presented in Table 1.

According to T-test, values in Mazandarani and Caucasians were not significantly different except for in soft tissue facial convexity, nose prominence, nasolabial angle, upper lip thickness, lower lip to E line and H line and soft tissue chin thickness ( $p < 0.05$ ).

Table 1.Means and standard deviations of soft tissue parameters

Soft tissue parameters	Mean	Standard deviation	Max	Min	Number
Facial convexity	157.9100	5.6409	180.00	146.00	100
Nose prominence	17.8600	3.2004	2900	11.00	100
Nasolabial angle	104.8900	13.3613	139.00	70.00	100
lower vertical height	69.1200	7.0758	89.00	51.00	100
Upper lip length	20.0900	3.4615	29.00	11.00	100
Upper lip thickness	12.8200	2.4137	20.00	7.00	100
Maxillary incisor exposure	4.1150	1.8988	9.00	1.00	100
Interlabial gap	2.4700	0.6506	4.00	1.00	100
Lower lip – menton height	46.1500	5.8592	64.00	30.00	100
Lower lip thickness	14.4300	2.2574	20.00	8.00	100
Lower lip protrusion	2.5400	1.5467	6.00	-2.00	100
Sub nasal perpendicular to Chin	3.1950	1.5191	9.00	0.00	100
Lower face throat angle	117.7600	9.5305	137.00	86.00	100
Throat length	61.5600	10.9104	85.00	28.00	100
Lower lip to E line	2.2500	2.0871	11.00	-2.00	100
Superior sulcus depth	2.3150	0.8308	5.00	0.00	100
Soft tissue sub nasal to H line	9.5900	2.5824	15.00	2.00	100
H angle	7.6800	1.0999	9.00	5.00	100
Lower lip to H line	6.4900	2.4225	12.00	2.00	100
Chin thickness	12.0400	2.6777	20.00	4.00	100
Middle to lower third	0.7904	0.1117	1.10	0.50	100
Sub nasal- stomion to stomion- menton	0.6796	0.1291	1.02	0.39	100
Vertical height ratio	1.1554	0.1672	1.67	0.67	100
Lower vertical height depth ratio	1.1475	0.1897	1.68	0.81	100
Vertical lip Chin ratio	0.4751	0.2888	0.78	0.32	100

## Discussion

Facial esthetics is based on both soft and hard tissues. Soft tissue features such as thickness, length, vertical and horizontal positions and the angles among constructed planes on soft tissue are measurable. The purpose of the present study was to evaluate the soft tissue proportions of normal facial profile

The mean of facial convexity according to Legan-Burstone analysis in Iranian population was 157.91° lower than Caucasian [6] and Japanese norms [8] and expressing more convex profile in north Iranian population. Nose prominence in Holdaway analysis was 17.86 millimeters lower than Caucasian norm. [4, 5] Nose prominence was greater in this study than the norms in Japanese [6], Turkish [16], and Saudi Arabian populations. [8] Nasolabial angle in this research was more obtuse than Caucasians [6] and Brazilian soft tissue norms [9], but was lower than Japanese norm. [7] The differences among these studies indicate the racial difference.

Upper lip length in this research was lower than the norms of Japanese [7] and Turkish population [16] but had no significant difference with Caucasians. [15] Upper lip thickness was lower than norms of Caucasian [6], and Japanese populations. In Mazandaran, people have thinner upper lip, but there was no significant difference between north Iranian population and Caucasians in terms of lip length. The distance of lower lip to E line was more this research than Caucasians [4, 5] and Japanese adults. [7] In addition, the distance of lower lip to H line was more than Caucasians, [4, 5] indicating more prominent lower lip in north Iranian population.

Thickness of chin soft tissue was more in mazandaranians than Caucasians [4, 5] but lower than norm in Japan. [7] Because the number of males and females was not the same, the comparison of the soft tissue norms between sexes was impossible; therefore, it

is suggested to compare the norms between two sexes in further studies.

## Conclusion

In comparison with Caucasians, people in Mazandaran have a more convex profile, greater nasolabial angle, more prominent lower lip, thicker chin soft tissue and shorter upper lip.

**Conflict of interest:** There is no conflict of interest.

## Authors' Contributions

The study was designed by Manouchehr Rahmatikamel and Valiollah Arash. The study data were collected by Abbasali Ostadrahimi. Analysis and interpretation of data, drafting of the manuscript and critical revision of the manuscript for important intellectual content were performed by Abbasali Ostadrahimi and Reza Ghorbanipour. Study supervision was conducted by Manouchehr Rahmatikamel.

## References

1. Bishara SE, Jakobsen JR, Hession TJ, Treder JE. Soft tissue profile changes from 5 to 45 years of age. *Am J Orthod Dentofacial Orthop* 1998; 114: 698–706.
2. Jacobson A, Jacobson RL, editors. Radiographic cephalometry : from basics to 3-D imaging. 2<sup>nd</sup> ed. Chicago: Quintessence Pub; 2006. p. 219-31.
3. Ioi H, Shimomura T, Nakata SH, Nakasima A, Counts AL. Comparison of anteroposterior lip positions of the most-favored facial profile of korean and japanese people. *Am J Orthod Dentofacial Orthop* 2008; 134: 490-5.
4. Holdaway RA. A soft-tissue cephalometric analysis and its use in orthodontic treatment planning. Part I. *Am J Orthod* 1983; 84:1-28.
5. Holdaway RA. A soft-tissue cephalometric analysis and its use in orthodontic treatment planning. Part II. *Am J Orthod* 1984; 85:279-93.
6. Bergman RT. Cephalometric soft tissue facial analysis. *Am J Orthod Dentofacial Orthop* 1999; 116:373-89.
7. Alcalde RE, Jinno T, Orsini MG, Sasaki A, Sugiyama RM, Matsumura T. Soft tissue cephalometric norms in japanese adults. *Am J Orthod Dentofacial Orthop* 2000; 118:84-9.
8. Kundi I. Cephalometric Soft tissue standard and gender dimorphism in nasal prominence estimated by holdaway's analysis in patients visiting college of dentistry, aljouw university. *J Contemp Dent Pract* 2017; 18:152-5.
9. Magnani MB, Nouer DF, Nouer PR, Pereira Neto JS, Garbui IU, Böeck EM. Assessment of the nasolabial angle in young brazilian black subjects with normal occlusion. *Braz Oral Res* 2004; 18:233-7.
10. Basciftci FA, Uysal T, Buyukerkmen A. Determination of holdaway soft tissue norms in anatolian turkish adults. *Am J Orthod Dentofacial Orthop* 2003; 123:395-400.
11. Ioi H, Nakata SH, Nakasima A, Counts AL. Comparison of cephalometric norms between Japanese and caucasian adults in antero-posterior and vertical dimension. *Eur J Orthod* 2007; 29: 493-9.
12. Ghorbany Javadpour F, Khane masjedi M. Evaluation soft tissue profile of face and anteroposterior lip position in persian adults and comparison with european-american standards. *Beheshti Univ Dent J* 2014; 32: 27-33. [In Persian]
13. Moghbel, B, Sodagar A, Etezadi T. Determination of Holdaway soft tissue norms in Iranian adolescents. *Iran J Orthod* 2010; 5: 32-8.
14. Legan HL, Burstone CJ. Soft tissue cephalometric analysis for orthognathic surgery. *J Oral Surg* 1980; 38:744-51.
15. Epker BN, Stella JP, Fish LC. Dentofacial deformities: Integrated orthodontic and surgical correction. 2<sup>nd</sup> ed. St Louis: Mosby Co, 1998; p.654.
16. Uysal T, Baysal A, Yagci A, Sigler LM, McNamara JA Jr. Ethnic differences in the soft tissue profiles of turkish and european-american young adults with normal occlusions and well-balanced faces. *Eur J Orthod*. 2012; 34:296-301.